

Bowel Diary

Date: I woke up at: I went to sleep at:

| Time | Food/drink consumed, medication (type + amount) | (√) For each bowel movement | Bowel urgency? Rate 1 mild - 3 strong | Any pains or discomfort? Rate 1 mild - 3 strong | Stool type (use the Bristol Stool Form Scale on next page) | Record any accidents/ leakage |
|------|--|--------------------------------------|--|---|---|-------------------------------------|
| 12am | | | | | | |
| 1am | | | | | | |
| 2am | | | | | | |
| 3am | | | | | | |
| 4am | | | | | | |
| 5am | | | | | | |
| 6am | | | | | | |
| 7am | | | | | | |
| 8am | | | | | | |
| 9am | | | | | | |
| 10am | | | | | | |
| 11am | | | | | | |
| 12pm | | | | | | |
| 1pm | | | | | | |
| 2pm | | | | | | |
| 3pm | | | | | | |
| 4pm | | | | | | |

| Time | Food/drink consumed, medication (type + amount) | (√) For each bowel movement | Bowel urgency? Rate 1 mild - 3 strong | Any pains or discomfort? Rate 1 mild - 3 strong | Stool type (use the Bristol Stool Form Scale on next page) | Record any accidents/ leakage |
|------|--|--------------------------------------|--|--|---|-------------------------------------|
| 5pm | | | | | | |
| 6pm | | | | | | |
| 7pm | | | | | | |
| 8pm | | | | | | |
| 9pm | | | | | | |
| 10pm | | | | | | |
| 11pm | | | | | | |

Bristol Stool Form Scale

Since it can be hard to state what is normal and what is abnormal, some health professionals use a scale to classify the type of stool passed. This helps assess how long the stool has spent in the bowel.

Type 1 has spent the longest time in the bowel and type 7 the least time. A normal stool should be a type 3 or 4 and depending on the normal bowel habits of the individual, should be passed once every one to three days.